

Evidence for Dyonic Structure of $SU(2)$ Lattice Gauge Fields Below T_c ¹

E.-M. Ilgenfritz^{a2}, B. V. Martemyanov^b, M. Müller-Preussker^a, S. Shcheredin^a
and A. I. Veselov^b

^a *Institut für Physik, Humboldt-Universität zu Berlin, D-10115 Berlin, Germany*

^b *ITEP, B. Cheremushkinskaya 25, Moscow, 117259, Russia*

Abstract

We report the observation of calorons with nontrivial holonomy and fractional topological charge objects in cooled lattice samples derived from $SU(2)$ equilibrium ensembles at $T < T_c$.

Characteristic features of QCD are related to the existence of lumps of topological charge in the vacuum. Usually, they are described as instantons or, at $T \neq 0$, calorons [1] and considered to be the agents of chiral symmetry breaking. Recently, this view has been questioned from different sides. At first, an old companion of instantons, the meron picture (see *e.g.* [2]) has been revived. One motivation is to deduce confinement from the dissociation of instantons into constituents. The semiclassical picture of instantons has been totally challenged by Horvath et al. [3]. Van Baal et al. [4] have stressed the existence of more general instanton solutions at $T \neq 0$, distinguished by nontrivial holonomy boundary conditions. A semiclassical approach to deal with this situation has been outlined for four dimensional $\mathcal{N} = 1$ supersymmetric gluodynamics [5].

The present talk reports on recent lattice evidence [6] for the existence of dyonic lumps in the confinement phase. They are expected to be constituents of the new caloron solutions and show the required color correlations. The new (KvB) caloron [4] is a finite-temperature (periodic) solution of the Euclidean field equations. It is selfdual with topological charge $Q = \pm 1$. The holonomy at asymptotics is not restricted to trivial values, $P = \exp \left(i \int_0^{1/T} A_4 dx_4 \right) \rightarrow P_\infty \notin Z(N_c)$. Depending on P_∞ , part of the parameter space corresponds to solutions with action (and topological charge) concentrated in N_c well-separated static constituents. One of the constituents is carrying the fermion zero eigenmode required to exist for a $Q = \pm 1$ object according to the Atiyah-Singer index theorem.

¹Work supported by DFG. B. V. M. and A. I. V. acknowledge receiving grants RFBR 02-02-17308 and 01-02-17456, INTAS 00-00111 and the CRDF award RP1-2364-MO-02.

²Talk presented by E.-M. I. at the 5th International Conference on Quark Confinement and the Hadron Spectrum, Gargnano, Italy, 10-14 Sep 2002

Monte Carlo configurations taken from confined and unconfined ensembles, respectively, have been "cooled" (by gradual minimization of action). We wanted to show [6] that nearly classical configurations of KvB type really pop up as background fields starting from the confinement phase. In fact, this happens at temperature close to T_c . In contrast, the deconfined phase does not support topologically nontrivial background fields. Monitoring the relaxation $\frac{dA}{d\tau} \sim -\frac{\partial S}{\partial A}$, cooling was stopped triggered by the gradient becoming *minimal within some window of action*, and snapshots of configurations were taken. On plateaux at moderate multiples of $S_{inst} = 2\pi^2$, the multi-lump structure is still comprehensible and can be analysed as described in the following. Figs. 1 - 3 show different portraits of the same selfdual *dyon-dyon pair* obtained by cooling under periodic gluonic boundary conditions with $S \approx 2\pi^2$. Almost static dyons are localizable by peaks of the topological density $q(x) > 0$ and the Polyakov loop $L(\vec{x}) \approx \pm 1$.

A complementary view is given by the spectrum of the (non-Hermitean) Wilson fermion Dirac operator. Real eigenvalues refer to a fermionic zero mode which, depending on periodic (p.b.c.) or antiperiodic (a.b.c.) fermion boundary conditions in time, is localized at one or the other constituent. Cooled configurations like this can be well fitted by the shape functions of analytic KvB solutions. Nonstatic $Q = 1$ caloron configurations can also be classified according to the KvB solutions. Unexpectedly, also *dyon-antidyon pairs* (with opposite topological charge but same sign Polyakov loop) have been detected in the same window of action and found to be almost stable. In this case a pair of almost-real modes exists, similar to those of an instanton-antiinstanton pair. The saddle point configurations in higher windows of action (and their cascade decay under cooling) can be interpreted in terms of such dyonic constituents.

Further details can be found in Ref.[6]. Corroborating the point of view dealt with here, *fermionic* results concerning *uncooled* configurations of $SU(3)$ gluodynamics have been reported [7, 8] after the conference.




Figure 1: Topological density $q(x)$ (left) and Polyakov loop $L(\vec{x})$ (right) of a dyon-dyon pair




Figure 2: Lowest fermionic eigenvalues for periodic (left) and antiperiodic (right) boundary conditions in Euclidean time




Figure 3: Localized fermionic zero mode, for periodic (left) and antiperiodic (right) boundary conditions in Euclidean time

References

- [1] D. J. Gross, R. D. Pisarski, L. G. Yaffe, Rev. Mod. Phys. **53** (1981) 43.
- [2] A. Montero and J. W. Negele, Phys. Lett. **B 533** (2002) 322.
- [3] I. Horvath et al., these proceedings and references therein.
- [4] T. C. Kraan and P. van Baal, Phys. Lett. **B 435** (1998) 389, Nucl. Phys. **B 533** (1998) 627.

- [5] N. M. Davies et al., Nucl. Phys. **B 559** (1999) 123.
- [6] E.-M. Ilgenfritz et al., Phys. Rev. **D 66** (2002) 074503.
- [7] C. Gattringer, e-print hep-lat/0210001, Phys. Rev. **D** in print.
- [8] C. Gattringer and S. Schaefer, e-print hep-lat/0212029.